

Health

6.6 Poorly children

- If a child appears unwell during the day, for example has a raised temperature, sickness, diarrhoea or pains, particularly in the head or stomach then the setting manager calls the parents and asks them to collect the child or send a known carer to collect on their behalf.
- If a child has a raised temperature, they are kept cool by removing top clothing, sponging their heads with cool water and kept away from draughts.
- A child's temperature is taken and checked regularly, using Fever Scans or other means i.e. ear thermometer.
- In an emergency an ambulance is called and the parents are informed.
- Parents are advised to seek medical advice before returning them to the setting; the setting can refuse admittance to children who have a raised temperature, sickness and diarrhoea or a contagious infection or disease.
- Where children have been prescribed antibiotics for an infectious illness or complaint, parents are asked to keep them at home for 48 hours.
- After diarrhoea or vomiting, parents are asked to keep children home for 48 hours following the last episode.
- Some activities such as sand and water play and self-serve snack will be suspended for the duration of any outbreak.
- The setting has information about excludable diseases and exclusion times.
- The setting manager notifies their line manager if there is an outbreak of an infection (affects more than 3-4 children) and keeps a record of the numbers and duration of each event.
- The setting manager has a list of notifiable diseases and contacts Public Health England (PHE) and Ofsted in the event of an outbreak.
- If staff suspect that a child who falls ill whilst in their care is suffering from a serious disease that may have been contracted abroad such as Ebola, immediate medical assessment is required. The setting manager or deputy calls NHS111 and informs parents.

HIV/AIDS procedure

HIV virus, like other viruses such as Hepatitis, (A, B and C), are spread through body fluids. Hygiene precautions for dealing with body fluids are the same for all children and adults.

- Single use vinyl gloves and aprons are worn when changing children's nappies, pants and clothing that are soiled with blood, urine, faeces or vomit.
- Protective rubber gloves are used for cleaning/sludging clothing after changing.
- Soiled clothing is rinsed and bagged for parents to collect.

- Spills of blood, urine, faeces or vomit are cleared using mild disinfectant solution and mops; cloths used are disposed of with clinical waste.
- Tables and other furniture or toys affected by blood, urine, faeces or vomit are cleaned using a disinfectant.

Nits and head lice

- Nits and head lice are not an excludable condition; although in exceptional cases parents may be asked to keep the child away from the setting until the infestation has cleared.
- On identifying cases of head lice, all parents are informed and asked to treat their child and all the family, using current recommended treatments methods if they are found.

****Paracetamol based medicines (e.g. Calpol)**

The use of paracetamol-based medicine may not be agreed in all cases. A setting cannot take bottles of non-prescription medicine from parents to hold on a 'just in case' basis, unless there is an immediate reason for doing so.

Further guidance

Guidance on infection control in schools and other childcare settings (Public Health Agency)

<https://www.publichealth.hscni.net/sites/default/files/>

[Guidance_on_infection_control_in%20schools_poster.pdf](#)

This policy was reviewed & updated on: 23/09/21 **and adopted by:** Discovery Vine Ltd

Date to be reviewed: September 2022 (or earlier if necessary)

Signed on behalf of Discovery Vine Ltd: *Lynda Garbutt*

Name of signatory: Lynda Garbutt

Role of signatory: Managing Director